



TRAVEL HEALTH PRICE LIST & SCHEDULE



Condition	Cost	Schedule
Altitude sickness	Dependent on supply	1 day prior to ascent, continued for at least 2 days once acclimatised, or once descent has begun
Chikungunya	£200	1 dose 3 weeks before travel
Cholera (oral)*	£75 per course	1 dose 10 days before travel OR 2 doses within 6 weeks of each other with second dose at least 1 week before travel
Hepatitis A*	£65	1 dose at least 2 weeks before travel with booster after 6 months
Hepatitis A&B	£85 per dose £170-£255 per course	2 doses: 0 then 6-12 months OR 3 doses: 0, 1, and 6 months OR rapid schedule (exceptional circumstances only): day 0, 7 and 21, prior to travel with booster 12 months later
Hepatitis B	£50 per dose £150 per course	3 doses: 0, 1 and 6 months OR 0, 1 and 2 months with booster 12 months later (exceptional circumstances)
Japanese Encephalitis	£115 per dose £230 per course	2 doses: day 0 and 28 OR rapid schedule (exceptional circumstances): day 0 and 7; final dose at least 1 week before travel
Malaria	Dependent on supply	1-2 days before entering a malaria-endemic area, continued during the period of exposure and for 4 weeks after leaving
	Dependent on supply	1-2 days before entering a malaria-endemic area, continued during the period of exposure and for 7 days after leaving
	Dependent on supply	10 days before departure, then once a week during period of exposure and for 4 weeks after leaving
Measles, Mumps & Rubella (MMR)	£45 per dose £90 per course	2 doses at least a month apart; final dose at least 2 weeks before travel
Meningitis ACWY	£60	1 dose at least 2 weeks before travel
Polio, Diphtheria, Tetanus*	£40	1 dose (booster) if >10 years since vaccination, at least 2 weeks before travel
Rabies	£80 per dose £240 per course	3 doses: day 0, 7 and 28 OR rapid schedule (exceptional circumstances): day 0, 3 and 7 with a booster at 12 months if needed; final dose at least 1 week before travel
Tick-borne Encephalitis	£65 per dose £195 per course	3 doses: 1 st injection day 0, 2 nd 1-3 months later; 3 rd 5-12 months after 2 nd injection OR day 0, day 14 and 5-12 months after 2 nd injection (special cases). Immunity after 2 doses: 2 nd dose at least 2 weeks before travel
Typhoid*	£40	1 dose at least 2 weeks before travel
Yellow Fever	£100	1 dose at least 10 days before travel

*these may be available on the NHS - check with your local surgery

[Book & fill out a pre-consultation form here](#)